

Maxillary sinusitis of odontogenic origin in relation to ectopic 2.8 associated with follicular dentigerous cyst: Combined transnasal and oral endoscopic approach, a case report

Mariagrazia Boccuto¹, Rosario Marchese Ragona² and Christian Bacci^{1*}

¹Unit of Oral Medicine, Pathology and Surgery, Section of Clinical Dentistry, University of Padova, Italy

²Department of Neurosciences, Section of Otolaryngology, University of Padova, Italy

Abstract

Purpose of the study: To describe the management of orosinus pathology by combined transnasal endoscopy and oral combined.

Methods: The 54-year-old patient underwent a dental and otolaryngological evaluation for left odontogenic maxillary sinusitis in relation to plausible dental follicular cyst of ectopic element 2.8. The symptoms reported at the time of access to the hospital were nasal obstruction and nocturnal rhonchopathy.

To the ENT evaluation in videorinoscopy with rigid optics, complex deviation of the nasal septum was relevated with not any evident formations or pathological secretions. While on inspection of the oral cavity the mucous membranes appeared unscathed. Alveolus of 1.6 previously extracted was evident.

The radiological examination, facial CT, revealed the left maxillary sinus almost completely occupied by a cystic appearance, with thin calcified walls and homogeneous content that has a dental element, probably the 2.8, which fenestrates the vestibular cortex of the lateral wall of the maxillary sinus. This lesion erodes the medial wall of the sinus, obliterating the ostio-meatal complex and imprinting the ipsilateral ethmoidal cells.

Biohumoral tests showed normal coagulation parameters, indices of renal function, liver and ionemia.

The patient under general anesthesia and oral intubation with a combined intervention of the left anterior FESS, intrasulcular flap from dental elements 2.7 to 2.3 with mesial releasing incision, moderate osteotomy, ectopic 2.8 extraction and enucleation of the cystic lesion with simultaneous closure of the orosinus communication with advancement of the Bichat adipose bolla and closure by first intention.

In the same session, the ENT moment is carried out trans nasally for total left uncinectomy, medium antrostomy with the union of the natural ostium and the accessory ostium. Bilateral lower turbinoplasty with bipolar forceps.

The patient was then controlled after 15 days and six months, showing good healing and no signs of recurrence at the rhinoscopic check on the physical examination of the oral cavity.

Results: based on the clinical and radiological aspect, the diagnosis of a follicular dentigerous cyst (WHO 2017) covered by a multi-layered non-keratinized paving epithelium, with moderate chronic inflammation, including gigantocellular and cholesteric crystals, is reached from the microbiological and histological examination. Necrotic amorphous material coexists including rare hyphae and fungal spores, with therefore mycotic and actinomycotic super infection.

Conclusions: The combined oral and nasal intervention, allowed by the collaboration between the oral surgeon and ENT, has made it possible to shorten the healing time and to resolve the pathology without any sign of recurrence.

Introduction

The physiological functions of the paranasal sinuses depend on the patency of the ostio-meatal complexes and the mucociliary functionality. The interruption of this equilibrium can lead to sinus pathology such as sinusitis. For a long time, it was believed that primary inflammation of the maxillary epithelium was the main cause of sinusitis; however, it was later shown that most diseases start from a deficiency of the functionality of the ostio-meatal complex which interrupts normal drainage and decreases ventilation inside the sinus itself. Localized sinus infections are

less common and can occur from focal areas of inflammation within the individual sinus: such as starting from a dental pathology.

10% of cases of maxillary sinusitis have an odontogenic origin: from dental infection, from dental trauma or from iatrogenic causes such as extractions, osteotomies or endosseous implants. In such cases, therapy requires the resolution of the odontogenic pathology in addition to the sinusal one [1].

The follicular cyst or dentigerous cyst (according to the most recent WHO classification of 2017 that reported this term in use)

develops around the crown of the included tooth and is caused by the accumulation of fluid between the reduced epithelium of the enamel and the coronal surface of the tooth.

Radiographically, unilocular, well defined radiolucency is observed from at the margin of the crown of an unerupted tooth. The affected tooth is often dislocated because of the cyst itself.

An inflammatory infiltrate consisting of lymphocytes, granulocytes and plasma cells is frequently associated.

As in root cysts, cholesterol crystals, hemosiderin deposits and Rushton bodies may be present [2].

The resolution of the pathology and the closure of the orosinusal communication, can be made use of different surgical techniques such as the use of flaps, grafts or other techniques [3].

Traditional mucoperiosteal flaps include sliding vestibular flap, rotated palatal flap, transposed palatal flap and lingual flaps. Other techniques use the combination of the aforementioned flaps, with possible advancement and isolation of the Bichat bolla and/or skin grafts.

- The grafts can be autogenous from the symphyseal area, retromolar area, zygomatic area or they can use membranes rich in fibrin, platelet gel and/or septal cartilage. The use of xenografts associated with the mobilization of mucoperiosteal flaps involves the use of freeze-dried porcine dermis, porcine connective membranes, bovine bone and guided tissue regeneration (GTR) techniques associated with bovine membranes. The allografts use instead of freeze-dried fibrin glue and GTR.

- Other techniques include third molar reimplantation, closure of the post-surgical site by primary intention, the use of metal plates or polymethyl methacrylate in association with palatal and buccal mucoperiosteal flaps, hydroxyapatite blocks and hemostatic gauze.

Clinical case

54-year-old man comes to our attention for left maxillary sinusopathy referring nasal obstructive symptoms especially on the left and nocturnal roncopathy with apneas.

The patient denies pain and asthma.

In anamnesis, he remembers previous surgery of appendectomy, chronic pathologies such as hypercholesterolaemia, for which he performs home therapy with Simvastatin 200 mg. Deny drug allergies.

Upon entering the Padova Teaching Hospital, he underwent a facial CT which shows the left maxillary sinus completely occupied by a cystic formation, with slender calcified walls and homogeneous content that includes a dental element, Ectopic 2.8 element.

This cystic formation medially obliterates the ostio-meatal complex and deforms the medial wall of the maxillary sinus causing a moderate impression on the ipsilateral ethmoid cells, even with partial obliteration of the left fronto-ethmoid recess, on the turbinates and on the nasal bone septum (Figures 1,2).

After this radiological assessment, ENT and dental clinical evaluations were performed: the first confirms the need to perform left and contextual septic plastic FESS (Functional Endoscopic Sinus Surgery) for complex deviation of the nasal septum highlighted following rhinoscopy and the second reiterates the need to close the orosinusal communication at the same time as the proposed act following enucleation of the cystic lesion and extraction of an ectopic 2.8 element.




Figure 1. Coronal Scan of CT




Figure 2. Sagittal Scan of CT

At intraoral physical examination, no alterations of the mucous membranes were present, the presence of fistulas or any pathological secretions.

Therefore, under general anesthesia, the oral procedure is first carried out with infiltration of local anesthetic. A Block of the left maxillary nerve is performed via descending palatal canal descending through the major palatal foramen, intraoral infraorbital nerve block of the middle and posterior superior alveolar nerve and paraperiosteal infiltration, with articaine with epinephrine 1: 200000 to provide deeper anesthesia by taking advantage of the high lipophilicity [4]

An intrasulcular incision from element 2.3 to element 2.7 and mesial releasing incision, full thickness flap and subsequent moderate osteotomy are performed, which allows the extraction of the ectopic element 2.8 and the enucleation of the cystic lesion, which highlights in the content hyperdense areas, compatible, clinically, with actinomycotic colonies.

The sample was sent for histopathological and microbiological analysis; the Bichat bulla [5] was advanced to occupies the space of communication itself, exploiting not only its volume but also its osteogenic capacities, due to the high presence of stem cells [6-9], subsequent closure by primary intention with 4/0 absorbable suture.

During the intranasal endoscopic moment, the left with a total uncinectomy [7] [Functional endoscopic sinus surgery, Slack 1998] and subsequent average antrostomy with union of the natural ostium of the maxillary sinus with the accessory ostium were performed. Follow with incision of the septal mucosa along the left condrovomerian ridge and subperichondral-subperiosteal detachment of the stenosing ridge, which was removed.

It ends with bilateral inferior turbinoplasty using bipolar forceps, positioning of a flexible silicone plate for each nasal fossa anchored to the septum with a suture point and a bilateral nasal swab.

At home, drug therapy was prescribed: anti-histamine rupatadine 10 mg for 30 days, nasal washes with physiological solution for 20 days, topical endonasal application of neomycin for 15 days and paracetamol 1000 mg [10] three time a day, for pain management.

The patient then returned to control after 15 days for the removal of the endonasal lamina showing good intraoral healing of the soft tissues, which allowed the removal of the previously positioned suture and after 6 months for rhinoscopic control which showed no sign of recurrence.

Results

Based on the clinical and radiological aspect, a microbiological investigation and microscopic examination leads to the diagnosis of a follicular dentigerous dental cyst coated with multi-layered non-keratinized paving epithelium, with moderate chronic inflammation, including gigantocellular and cholesteryl crystals. Necrotic amorphous material coexists including rare hyphae and fungal spores, with actinomycotic superinfection.

Conclusions

The combined oral and nasal intervention, allowed by the collaboration between the oral surgeon and ENT, has made it possible to shorten the healing time and to resolve the pathology without any sign of recurrence.

Furthermore, according to our approach, the optimal management of the sinus pathology related to the odontogenic cause is no longer purely dental or otolaryngology when there is involvement of several paranasal sinuses and a complete or partial obliteration of the osteomeatal complex appears.

Therefore the interventional contemporaneity in order to exclude recurrences and resolve the pathology is desirable.

References

1. Sinusitis. In: Oral and Maxillofacial pathology (3rd Edn). Neville, Damm, Allen, Bouquot Eds: Saunders Elsevier Pbl: United States of America, 2009: 207-208
2. Cisti dentigera (follicolare). In: Manuale di patologia e medicina orale, terza edizione. Giuseppe Ficarra Eds: McGraw-Hill Pbl: Milano, 2006: 353-355.
3. Kiran Kumar Krishanappa S, Prashanti E, Sumanth KN, Naresh S, Moe S, et al. Interventions for treating oro-antral communications and fistulae due to dental procedures. Cochrane Database Syst Rev. 2016; CD011784.
4. Yapp KE, Hopcraft MS, Parashos P. Articaine: a review of the literature. Br Dent J. 2011; 210: 323-329.
5. Horowitz G, Koren I, Carmel NN, Balaban S, Abu-Ghanem S, et al. One stage combined endoscopic and per-oral buccal fat pad approach for large oro-antral-fistula closure with secondary chronic maxillary sinusitis. Eur Arch Otorhinolaryngol. 2016; 273: 905-909.
6. Salehi-Nik N, Rad MR, Kheiri L, Nazeman P, Nadjmi N, et al. Buccal fat pad as a potential source of stem cells for bone regeneration: a literature review. Hindawi Stem Cells International. 2017: 8354640.
7. Robert Slack, Grant Bates. Functional endoscopic sinus surgery. American Family Physician. 1998; 58: 707-718.
8. Bacci C, Lucchiari N, Valente M, Della Barbera M, Frigo AC, et al. Intra-oral bone harvesting: two methods compared using histological and histomorphometric assessments. Clin Oral Implants Res. 2011; 22: 600-605.
9. Bacci C, Sivoilella S, Brunello G, Stellini E. Maxillary sinus bone lid with pedicled bone flap for foreign body removal: the piezoelectric device. Br J Oral Maxillofac Surg. 2014; 52: 987-989.
10. Manani G, Bacci C, Zanette G, Facco E. Contemporary state of sedation in dentistry. Dental Cadmos. 2012; 80: 357-369.

***Correspondence:** Christian Bacci, Department of Neurosciences, Section of Otolaryngology, University of Padova, Italy, E-mail: christian.bacci@unipd.it

Rec: Jan 31, 2020; Acc: Feb 12, 2020; Pub: Feb 15, 2020

Global Dentistry. 2020;3(1):131
DOI: gsl.jgd.2020.000131

Copyright © 2020 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY).